

**TABLE 1. Population Demographics**

	Entire Cohort (n=181)	HIV + with Lipodystrophy (n=92)	HIV+ without Lipodystrophy (n=89)	p-value between Lipodystrophy groups
Age (years)*	43 (38, 50)	44 (38, 50)	41 (36, 46)	0.03 <sup>¥</sup>
Race	32% Hispanic, 9% black, 52% white, 2% Asian, 5% other	34% 9% 52% 2% 3%	29% 10% 52% 2% 7%	0.83 <sup>Φ</sup>
Sex (Male: Female)	128: 53	60: 32	68: 21	0.10 <sup>Φ</sup>
CD4 count (cells/ $\mu$ L)*	463 (262, 650)	496 (317, 678)	421 (242, 641)	0.06 <sup>¥</sup>
Log <sub>10</sub> HIV viral load (copies/mL)*	2.6 (2.6, 3.4)	2.6 (2.6, 2.6)	2.6 (2.6, 3.8)	0.04 <sup>¥</sup>
Lipodystrophy Assessment Score*	1.5 (0, 3.0)	3.0 (2.0, 4.5)	0.5 (0, 1.0)	<0.0001 <sup>¥</sup>
SIBID-S score*		2.1 (1.4, 2.9)	1.3 (0.6, 2.0)	<0.0001 <sup>¥</sup>
BIQLI score*		-0.5 (-1.2, 0.4)	0.4 (-0.4, 1.5)	<0.0001 <sup>¥</sup>
Hypertension (N (%))	77 (43%)	43 (47%)	34 (38%)	0.25 <sup>Φ</sup>
Diabetes (N (%))	36 (20%)	20 (22%)	16 (18%)	0.53 <sup>Φ</sup>
Dyslipidemia (N (%))	102 (56%)	59 (64%)	43 (48%)	0.03 <sup>Φ</sup>
HCV (N (%))	22 (12%)	13 (14%)	9 (10%)	0.41 <sup>Φ</sup>
AIDS (N (%))	102 (56%)	59 (64%)	43 (48%)	0.03 <sup>Φ</sup>

\*Expressed as median (interquartile range).

<sup>¥</sup>p-value by Wilcoxon rank sum test. <sup>Φ</sup>p-value by chi-square statistics.