

Table 2. Healthcare utilization outcomes according to lipodystrophy status.

Healthcare utilization variable	HIV + with Lipodystrophy (N=92)	HIV+ without Lipodystrophy (N=89)	p-value
Total Healthcare Encounters	19 (10, 26)	11 (7, 20)	0.0002
Total Healthcare Costs	\$4,731 (\$2,427, \$8,242)	\$3,013 (\$1,493, \$5,047)	0.007
Clinic Visits	18 (9, 26)	10 (6, 19)	0.0003
Clinic Costs	\$2,841 (\$1,274, \$4,478)	\$2,193 (\$939, \$3,653)	0.06
Emergency Room Visits	0 (0, 1)	0 (0, 0)	0.20
Emergency Room Costs	\$0 (\$0, \$469)	\$0 (\$0, \$0)	0.20
Hospital Admission during 12 month period (%)	21%	9%	0.03[¥]
Length of Stay (days)*	3 (2, 8)	3 (1, 5)	0.21
Hospital Admission Costs*	\$4,668 (\$0, \$8,163)	\$4,775 (\$1,608, \$9,508)	0.67

Data reported as median (IQR) unless otherwise specified. p-value by Wilcoxon rank sum test unless otherwise specified.

[¥]p-value by chi-square analysis. *Data from hospitalized patients only.